(leading to M_{top} Measurement) Z + Jets Study 8.4.0 Update


Jimmy Proudfoot Sept 14,2004 _


Event Sample


- Files:
 - /castor/cern.ch/user/a/agupta/zeemm/*
 - (i.e. DC1 simulation, with 8.4.0 reconstruction)
- Selected good electrons
 - if(nt.eg_lsEM[i] == 0 && nt.eg_trkmatchnt_X[i] >0)
 - 0.7< E/p < 1.3
 - Required 2 oppositely charged electrons
- Mass of electron pair 71.0 < M_{ee} < 111.0 GeV
- Gives about 800 events to use to study jet balancing
- Use (η,φ) match to identify the jets which contain the electrons

Variables Studied so Far

```
Pt(Z)
Pt(Leading Jet)
\Delta \phi between reconstructed Z and leading jet
\Delta R between leading jet and closest one in
  (R, \phi)
Et of the closest jet
M(Z)
E/p for electrons
Number of jets in the event
```








Fitted Zee mass is 88.58GeV


Why is it low?


Is it an error in the way I extract the electron 4-vectors from CBNT?


Does it assume a different calibration from that needed to establish the precision EM scale?


Does it reflect the bremmstrahlung seen in the E/p distribution?











Observations & Work in Progress

- Need to better understand the jet Et and multiplicity distributions.
 - Note that almost EVERY event has a second jet with Et>10GeV, and about half have a jet with Et>30GeV
- Why is the Z mass low by ~3%
- Need to go back and verify that I am correctly interpreting the numbers in the CBNT ntuple
- Need to make some appropriate cuts on the second jet and compare my results to those obtained earlier
- Start similar checkout on DC2 files